

Health Care Quality: A Consumer Perspective

Nancy S. Green, MD
Medical Director
March of Dimes



Associate Professor, Pediatrics
Assistant Professor, Cell Biology
Albert Einstein College of Medicine
April 2003

The “Well Child Health Care” Syndrome*

- Trust that children will turn out OK
- Lack of clear mechanisms for
 - Obtaining data
 - Making decisions - Taking responsibility

Private terminology, NS Green

The Story's Beginning

- Child born full term in 1995
- Uncomplicated delivery of a healthy newborn

Day 1 of Life

- 23 hours of age: nurse in newborn nursery noted jaundiced
- Jaundice not reported to the family or doctor
- Progression of jaundice not noticed
- No bilirubin or other tests

Day 2 of Life

- 33 hours of age: Jaundice continued but no bilirubin levels or other evaluation
- Normal physical exams and feeding
- 36 hours of age: Discharged home
- Pediatric follow up visit scheduled in 7 days

Parent Education

- The hospital's "parent education" consisted of:
 - a brochure
 - suggestion to put the baby in the window
and
 - assurance that jaundice was normal and the
baby was fine

1st Week of Life: Out-patient Problems

- Day 4 of life: began to breast feed with a weak suck and became lethargic.
- Mother called newborn nursery to tell them that he was "still yellow, lethargic and was feeding poorly"
- Mother reassured

More out-patient Problems

- Checked at visit to pediatrician's office
- Noted to look progressively jaundiced
- Bilirubin level not assayed

Re-hospitalized

- Hospitalized at 5 days of age
- Bilirubin was tested for the first time
- 35 mg/dcl = quite high
- Treatment limited to phototherapy
- Exchange transfusion was not done

More Problems

- Child's blood type never determined
- Admitting Pediatric resident incorrectly documented child's blood type - without typing done
- Blood incompatibility was "ruled out"

Signs of Neurologic Damage

- Day 6: developed respiratory distress
- Neurologic abnormalities (high pitched cry, increased tone and opisthotonos)
- Family not informed that this was typical Kernicterus.

False Reassurance

- Instead, reassured that Kernicterus “didn't happen any more in the United States”

More...


- Remained in the hospital for 5 days
- Several tests performed and family told that “none were significant”
- At discharge, child had:
 - multiple neurologic abnormalities
 - jaundice
 - poor feeding

Diagnosis

- 18 months of age - diagnosed with Kernicterus
- Age 2 years: child's blood type determined
- Retrospective diagnosis made of hyperbilirubinemia
 - caused by ABO incompatibility with isoimmunization

This is Kernicterus

- 7 years of age:
 - athetoid cerebral palsy
 - neurosensory hearing loss
 - dental dysplasia
 - ocular abnormalities
 - cannot walk independently or crawl
 - impaired speech
 - drooling



Sister – Lessons Learned

- Two and a half years later: child's sister born
- Parents had her cord blood sent for blood typing
- Confirmed that she was also a candidate for an ABO incompatibility

Correct Treatment

- 16 hours: she was noted to be jaundiced by the nursing staff
- Standing orders for a bilirubin test (per parental request)
- Bilirubin level was 13.5mg/dcl
 - (moderately high)
- Phototherapy initiated



A Normal Outcome

- Day 4: Sister discharged
- Discharged with home care nurse for 7 days to assess health and to follow blood bilirubin levels
- She is fine.

Fundamental Medical-Laboratory- Communication Problems

“Well Child Health Care Syndrome”

- Very common Pediatric problem
- Time-dependent
- Staff incredulity
- Inadequate or no Lab testing done

Multiple System Problems

- Inadequate staff observations
- Multiple lapses in follow-up of abnormalities
- Multiple medical staff involved
- In-patient and office lapses
- Insufficient use of laboratory to confirm clinical impressions

Interactions with Family

- Inadequate communication with family
- Inadequate parent education
- Inappropriate reassurance
- No corroborating laboratory and clinical data

Lessons Learned

- Multiple layers of error
- Lack of or inadequately enforced medical/nursing responsibilities
- Laboratory use: too late
- Unclear primary staff responsibility
- Family communication: too “rosy”

Prevention: Medical Needs

- Need routinized approach = Algorithm
 - e.g., possible mandatory Bilirubin screening in newborn nursery
- Mandatory provider participation
- Pediatric F/U = too late!

Prevention: Communication Needs

- Better parent educational materials
- Better attention to parental concerns
- Better use of laboratory data to address concerns